

Comparison of STI Screening Recommendations for Pregnant Women				
STI	USPSTF	CDC	AAFP	ACOG
Chlamydia	Screen women younger than 25 years and others at increased risk	Screen all	Screen women 25 years and younger and others at increased risk	Screen women at increased risk
Gonorrhea	Screen women younger than 25 years and others at increased risk	Screen women at increased risk	Screen women at increased risk	Screen women at increased risk
Syphilis	Screen all	Screen all	Screen all	Screen all
HIV	Screen all	Screen all	Screen all	Screen all
Hepatitis B	Screen all	Screen all	Screen all	Screen all
Hepatitis C	No specific recommendation	Screen women at increased risk	No specific recommendation	Screen women at increased risk
HSV	Do not screen	No specific recommendation	Do not screen	No specific recommendation
HPV*	No specific recommendation	No specific recommendation	No specific recommendation	No specific recommendation
<p>STI = sexually transmitted infection; USPSTF = U.S. Preventive Services Task Force; CDC = Centers for Disease Control and Prevention; AAFP = American Academy of Family Physicians; ACOG = American College of Obstetricians and Gynecologists; HIV = human immunodeficiency virus; HSV = herpes simplex virus; HPV = human papillomavirus; Pap = Papanicolaou</p> <p>* No treatment available; currently used to stratify risk of cervical neoplasia</p>				
<p>Meyers, D., Wolff, T., Gregory, K., et al. <i>USPSTF Recommendations for STI Screening</i>. Originally published in <i>Am Fam Physician</i> 2008;77:819-824. Agency for Healthcare Research and Quality, Rockville, MD. (http://www.ahrq.gov/clinic/uspstf08/methods/stinfections.htm)</p>				